PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :		(11) International Publication Number: WO 98/1519)5
A23L 1/302, 1/035, 2/52, A61K 9/00, 9/107	A1	(43) International Publication Date: 16 April 1998 (16.04.9	18)
(21) International Application Number: PCT/EP (22) International Filing Date: 2 October 1997 (6)		BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, M	
(30) Priority Data: 60/027,758 7 October 1996 (07.10.96) (71) Applicant: F.HOFFMANN-LA ROCHE AG [CH/C Grenzacherstrasse, Ch-4070 Basle (CH). (72) Inventors: CHEN, Chyi-Cheng; 26 Sequoia Place, W	СН]; 1:		
07470 (US). MIX, Jeffrey, Mark; 112 Summit Bloomfield, NJ 07003 (US). RAMANATHAN Jayanthan; 16H River Road, Nutley, NJ 07110 (US)	Avenu Srirar	e,	
(74) Agent: KELLENBERGER, Marcus; 124 Grenzach CH-4070 Basle (CH).	erstrass	е,	
• -			
(54) Title: VITAMIN PREPARATIONS FOR REVERAC	TE APE	LICATIONS	

(57) Abstract

The present invention provides an optically clear composition of fat soluble vitamins useful to fortify beverages. In addition, beverages which have been nutritionally supplemented using the optically clear composition of fat soluble vitamins are also provided for in the present invention.

BNSDOCID: <WO___ __9815195A1_I_>

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg .	SN	Senegal
ΑU	Australia	GA	Gabon	. LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
· BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	18	Iceland	MW	Malawi	US	United States of America
CA	Салада	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NSDOCID: <WO_____9815195A1_I_>

Vitamin Preparations for Beverage Applications

The present invention relates to improvements in nutritious vitamin and mineral supplemented beverages which contain vitamin E and other fat soluble vitamins. In particular, methods for preparing beverages are provided as are beverages containing fat soluble vitamins formulations.

Vitamin supplements for human and veterinary use are commonplace. Some diets, or heavy physical exercise, may require the intake of considerable quantities of vitamins apart from those generally obtained through what otherwise would be considered a balanced diet.

Vitamin supplementation is also important, primarily for those who have inadequate diets, including growing children. In Central and South American countries where the dietary intake of vitamins are low in the general population, such vitamin supplement would have great value.

Sport beverages, such as, for example GATORADE, and vitamin supplemented waters, are beverages where the addition of vitamins is of interest. Also of interest are beverages used to restore electrolytes lost through diarrhea, for example, Pedialyte. Additionally, carbonated beverages, such as, for example, flavored seltzer waters, soft drinks or mineral drinks, as well as non-carbonated fruit juices, fruit punches and concentrated forms of these beverages.

One object of the present invention is to provide an optically clear composition of fat soluble vitamins which are present in a nutritionally supplemental amount that can be added to beverages. Another object of the present invention is that the optically clear composition does not alter the sensory properties of the beverage to which it is added and that the composition is stable in the finished beverage (for example, no ringing or surface filming). Another object of the present invention is to provide a

15

10

5

20

25

BNSDOCID: <WO__

__9815195A1_l_>

15

20

25

30

35

beverage containing the optically clear composition of fat soluble vitamins which are present in a nutritionally supplemental amount.

Yet another object of the present invention is to provide an optically clear composition of vitamin E that can be added to beverages so that the amount of vitamin E is present in a nutritionally supplemental amount, typically from about 1 to about 30 milligrams/serving, serving being eight (8) ounces (or about 240 milliliters).

Yet another object of the present invention is to provide beverages and beverage concentrates which contain the optically clear composition of fat soluble vitamins which are present in a nutritionally supplemental amount.

Thus, the present invention relates to a vitamin supplement which comprises (A) (1) a fat soluble vitamin together with (2) an emulsifier which is then mixed into (B) a liquid to form an optically clear solution. The ratio of (1) to (2) is from about 1:1 to about 1:19 and the ratio of (A) to (B) is from about 1:0.5 to about 1:16. Preferably, the ratio of (1) to (2) is from about 1:1 to about 1:9, more preferably from about 1:1 to about 2:3, and most preferably about 2:3. Preferably, the fat soluble vitamin is selected from the group consisting of vitamin E or its esters, vitamin A or its esters, vitamin K, and vitamin D3 and mixtures thereof. More preferably, the fat soluble vitamin is vitamin E and its esters.

The emulsifier is preferably a non-ionic surfactant having good hydrophilic (strong affinity to water) and lipophilic (strong affinity to dispersed phase) characteristics. Additionally, the emulsifier should be GRAS (generally recognized as safe) or an approved material for food consumption as determined by the various regulatory agencies world wide. As an emulsifier, polyoxyethylene (20) sorbitan mono-oleate is a preferred material. It is also known as polysorbate 80 and is available commercially from, among other sources, ICI as Tween 80

As fat soluble vitamins, vitamin E or its esters (for example, vitamin E acetate), vitamin A or its esters (for example, vitamin A acetate and vitamin A palmitate), vitamin K (phytomenadione) and vitamin D3 (cholecalciferol) are

10

15

20

25

contemplated in the present invention. Also included are mixtures of said vitamins. Vitamin E or its esters is the preferred fat soluble vitamin, with vitamin-E acetate being most preferred.

The present invention is also directed to beverages containing the optically clear composition of fat soluble vitamins which are present in a nutritionally supplemental amount. Typical beverage or beverage concentrate compositions comprise at least about 0.05% by weight of fruit or cola flavor or at least about 3% by weight of fruit juice, a nutritionally supplemental amount of fat soluble vitamins added as an optically clear composition, water if necessary to bring the weight of the beverage to 100%, and optionally, a sweetener.

These beverages can be carbonated beverages e.g., flavored seltzer waters, soft drinks or mineral drinks, as well as non-carbonated fruit juices, fruit punches and concentrated forms of these beverages.

By way of example, the fruit juices and fruit flavors used herein include grape, pear, passion fruit, pineapple, banana or banana puree, apricot, orange, lemon, grapefruit, apple, cranberry, tomato, mango, papaya, lime, tangerine, cherry, raspberry, carrot and mixtures thereof.

Additionally, artificial flavors, e.g. cola, or natural flavors derived from these juices can be used in the beverages. Chocolate flavors and other non-fruit flavors can also be used to make beverages containing the vitamin and mineral supplement.

Beverages, especially juice and cola beverages, which are carbonated in the manner of soft drinks, as well as "still" beverages and also contemplated are nectars and full-strength beverages or beverage concentrates which contain at least about 45% by weight of juice.

Additionally, milk, obtained from cows or synthetic, is a contemplated beverage to which the optically clear composition of fat soluble vitamins can be added.

35

The liquid of (B) can be water, the beverage to be supplemented, as well as mixtures thereof.

As used herein, the term "fruit juice product" refers to both fruit juice beverages and fruit juice concentrates which comprise at least about 45% fruit juice.

Sport beverages are also contemplated as beverages which can be supplemented by the optically clear composition of fat soluble vitamins of the present invention. Typical sport beverages contain water, sucrose syrup, glucose-fructose syrup, and natural or artificial flavors. These beverages can also contain citric acid, sodium citrate, monopotassium phosphate, as well as other materials which are useful in replenishing electrolytes lost during perspiration.

15

10

5

As used herein, the term "fruit juice beverage" refers to a fruit juice product which is in a single-strength, ready-to-serve, drinkable form. Fruit juice beverages of the present invention can be of the "full-strength" type which typically comprise at least about 95% fruit juice.

20

Full strength fruit juice beverages also include those products of 100% fruit juice such as, for example, orange, apple, raspberry, cherry, apricot, pear, grapefruit, grape, lime, tangerine, carrot, pineapple, cranberry, tomato, and various mixtures thereof.

25

Fruit juice beverages also include extended juice products which are referred to as "nectars". These extended juice products typically comprise from about 50% to about 90% fruit juice, preferably, from about 50% to about 70% fruit juice. Nectars usually have added sugars or artificial sweeteners or carbohydrate substitutes.

30

35

As used herein, the term "fruit juice concentrate" refers to a fruit juice product which, when diluted with the appropriate amount of water, forms drinkable fruit juice beverages. Fruit juice concentrates within the scope of the present invention are typically formulated to provide drinkable beverages when diluted with 3 to 5 parts by weight water.

As used herein the term "beverage concentrate" or "beverage syrup" refers to a mixture of flavors, water and from about 10% to about 60% sugar or carbohydrate substitute, i.e. sucrose, dextrose, corn syrup solids, fructose, dextrins, polydextrose and mixtures thereof.

5

As used herein, the term "fruit juice materials" refers to concentrated fruit juice, plus other fruit juice materials such as fruit juice aroma and flavor volatiles, peel oils, and pulp or pomace.

10

15

٠, :

As used herein, the term "citrus juice" refers to fruit juices selected from orange juice, lemon juice, lime juice, grapefruit juice, tangerine juice and mixtures thereof.

The flavor component of the beverages and beverage concentrates contains flavors selected from fruit flavors, botanical flavors and mixtures thereof. As used herein, the term "fruit flavor" refers to those flavors derived from the edible reproductive part of a seed plant, especially one having a sweet pulp associated with the seed. Also included within the term "fruit flavor" are synthetically prepared flavors made to simulate fruit flavors derived from natural sources. Particularly preferred fruit flavors are the citrus flavors including orange, lemon, lime and grapefruit flavors. Besides citrus flavors, a variety of other fruit flavors can be used such as apple, grape, cherry, pineapple, mango and papaya flavors and the like. These fruit flavors can be derived from natural sources such as fruit juices and flavor oils, or can be synthetically prepared.

25

30

20

As used herein, the term "botanical flavor" refers to flavors derived from parts of a plant other than the fruit; i.e. derived from nuts, bark, roots and leaves. Also included within the term "botanical flavor" are synthetically prepared flavors made to simulate botanical flavors derived from natural sources. Examples of such flavors include cola, tea, and the like. Botanical flavors can be derived from natural sources such as essential oils and extracts, or can be synthetically prepared.

35

The flavor component can comprise a blend of various flavors, e.g. lemon and lime flavors, cola flavors and citrus flavors to form cola flavors, etc. If desired, fruit juices such as orange, lemon, lime, apple, grape and like juices

10

15

20

25

can be used in the flavor component. The flavors in the flavor component are sometimes formed into emulsion droplets which are then dispersed in the beverage concentrate. Because these droplets usually have a specific gravity less than that of water and would therefore form a separate phase, weighting agents (which can also act as clouding agents) are typically used to keep the emulsion droplets dispersed in the beverage. Examples of such weighting agents are brominated vegetable oils (BVO) and rosin esters, in particular the ester gums. See L.F. Green, Developments in Soft Drinks Technology, Vol. 1, (Applied Science Publishers Ltd. 1978), pp. 87-93, for a further description of the use of weighting and clouding agents in liquid beverages. Besides weighting agents, emulsifiers and emulsion stabilizers can be used to stabilize the emulsion droplets. Examples of such emulsifiers and emulsion stabilizers include the gums, pectins, celluloses, polysorbates, sorbitan esters and propylene glycol alginates. See L.F. Green, supra at p. 92.

The particular amount of the flavor component effective for imparting flavor characteristics to the beverages and beverage concentrates ("flavor enhancing") can depend upon the flavor(s) selected, the flavor impression desired, and the form of the flavor component. The flavor component can comprise at least 0.05% by weight of the beverage composition, and typically from 0.1% to 2% by weight for carbonated beverages. When fruit juices are used as the flavor, the flavor component can comprise, on a single-strength basis, up to 25% fruit juice by weight of the beverage, preferably from 5% to 15% fruit juice by weight for carbonated beverages.

Carbon dioxide can be introduced into the water which is mixed with the beverage syrup or into the drinkable beverage after dilution to achieve carbonation. The carbonated beverage can be placed into a container such as a bottle or can and then sealed. Any conventional carbonation methodology can be used to make the carbonated beverages of this invention.

30

35

The amount of carbon dioxide introduced into the beverage will depend upon the particular flavor system used and the amount of carbonation desired. Usually, carbonated beverages of the present invention contain from 1.0 to 4.5 volumes of carbon dioxide. The preferred carbonated beverages contain from 2 to about 3.5 volumes of carbon dioxide.

The present invention is also particularly suited for the supplementation of beverages and beverage concentrates, including citrus juices. The beverages can contain from 3% to 100% juice or from about 0.05% to about 10% of an artificial or natural flavor, particularly orange juice. The concentrated orange juice, orange juice aroma and flavor volatiles, pulp and peel oils used in the method of the present invention can be obtained from standard orange juice. See Nagy et al, Citrus Science and Technology, Volume 2, (AVI Publishing Co. 1977), pp 177-252 for standard processing of oranges, grapefruit and tangerines. (See also Nelson et al, Fruit and Vegetable Juice Processing Technology (3rd Ed., AVI Publishing 1980),pp. 180-505 for standard processing of noncitrus juices such as apple, grape, pineapple, etc. to provide sources of juice and juice materials for noncitrus juice products).

Juices from different sources are frequently blended to adjust the sugar to acid ratio of the juice. Different varieties of oranges can be blended or different fruit juices can be blended to get the desired flavor and sugar to acid ratio. A sugar to acid ratio of from about 8:1 to about 20:1 is considered acceptable for fruit juices. However, preferred sugar to acid ratios are typically from about 11:1 to about 15:1, especially for citrus juices

20

5

10

15

Sweeteners include the sugars normally present in fruit juice products, for example glucose, sucrose, and fructose. Sugars also include high fructose corn syrup, invert syrup, sugar alcohols, including sorbitol, refiners syrup, and mixtures thereof.

25

30

35

BNSDOCID: <WO_

__9815195A1_I_>

In addition to sugar, extended fruit juice beverages of the present invention can contain other sweeteners. Other suitable sweeteners include saccharin, cyclamates, acetosulfam, L-aspartyl-L-phenylalanine lower alkyl ester sweeteners (e.g. aspartame). A particularly preferred sweetener for use in such extended juice products is aspartame.

For single-strength fruit juice beverages, the sugar content can range from about 2° to about 16° Brix. Typically, the sugar content of such beverages depends upon the amount of fruit juice contained herein. For full-strength beverages containing at least about 95% fruit juice, the sugar content is typically from about 5° to about 14° Brix. For extended juice beverages which

10

15

20

25

30

35

comprise from about 50% to about 90% fruit juice, the sugar content is typically from about 5° to about 13° Brix (no other sweetener) or from about 2° to about 8° Brix (other sweetener containing). For fruit juice concentrates according to the present invention, the sugar content can range from about 6° to about 75° Brix. Typically, the sugar content of these juice concentrates is from about 20° to about 50° Brix. For orange juice concentrates, the sugar content is preferably from about 35° to about 50° Brix.

The amount of the sweetener effective in the beverages of the invention depends upon the particular sweetener used and the sweetness intensity desired. For noncaloric sweeteners, this amount varies depending upon the sweetness intensity of the particular sweetener. For sugar, this amount can be from 1% to 14% (typically from 6% to 14%) by weight for carbonated beverages. Preferred beverages contain from 9% to 13% by weight sugar. In determining the amount of sugar for beverages of the present invention, any sugar or other sweetener present in the flavor component, such as in fruit juice, is also included. Low-calorie sweetener combinations containing a noncaloric sweetener such as aspartame and a sugar such as high fructose corn syrup can also be used in beverages. For beverage syrups, the amount of sugar in a beverage syrup is from about 10% to about 60%, and preferably from about 40% to about 60%.

The various beverage and beverage concentrates can be packaged in conventional packages for the particular beverage or beverage concentrates which are nutritionally supplemented by the optically clear composition of fat soluble vitamins. In some instances, the concentrates are frozen.

The optically clear composition of the invention can be made by mixing the fat soluble vitamin and the emulsifier, in conventional mixing apparatus, at a temperature between about room temperature (~22°C) and about 90°C, preferably between about 35°C and 45°C, so as to form a homogeneous mixture. This homogeneous mixture is then allowed to cool to room temperature. The cooled mixture is then mixed with water at a temperature of from about 30°C to about 35°C, the ratio of the mixture to water being from about 1:0.5 to about 1:16. A preferred ratio range of mixture to water is from about 1:1 to about 1:9. When the composition and water are mixed, a gel-like

10

20

to fluid mixture is formed. While not being restricted to this theory, it is thought that the mixture has liquid crystal and liposome tendencies.

Additionally, instead of mixing the cooled mixture in water as described above, the cooled mixture could also be mixed into an appropriate amount of the beverage which is to be fortified in a like manner as described above. Such amount of beverage could then be used as a concentrate. Should the cooled mixture be added to water or to the beverage to be used as a concentrate, the resulting mixture should be pasteurized to avoid microbial spoilage. Such pasteurization methods are well known to those of ordinary skill in the art.

The following examples served to demonstrate the present invention.

Example 1 (control)

15 A Formulation:

Vitamin E acetate (4 grams) and Polysorbate 80 (polyoxyethylene (20) sorbitan mono-oleate; 6 grams) were mixed in a suitable container and heated to about 40°C to give a homogeneous mixture. The mixture was used for beverage fortification for the following beverages, Gatorade Lemon Ice and apple juice, as follows.

B. Fortification:

250 mg of the formulation A, containing 100 mg of vitamin E acetate, was dispersed in a sample of each of the aforementioned beverages to form a 100 g stock solution of each beverage. 1.25 g, 2.50 g or 3.75 g of the stock solution of each beverage was placed into a beaker. Into the respective beakers were placed 28.75 g, 27.50 g or 26.25 g of each of the respective beverages. Each
 30 beaker was stirred to disperse the materials therein to form beverages containing 1.25 mg, 2.50 mg and 3.75 mg of vitamin E acetate per 30 g of beverage. Turbidity of the beverages was measured using a nephelometer (Monitek, Model TA 1).

Example 2 (Inventive formulation)

WO 98/15195 PCT/EP97/05421 10

A Formulation:

Step 1.-Vitamin E acetate (4 grams) and Polysorbate 80 (polyoxyethylene (20) sorbitan mono-oleate; 6 grams) were mixed in a suitable container and heated to about 40°C to give a homogeneous mixture.

Step 2. The above mixture of Step 1(5 grams) and water (5 grams) were mixed and stirred at about 30-35°C until the mixture became gel-like. This mixture was used for beverage fortification for the following beverages, Gatorade Lemon Ice and apple juice, as follows.

B. Fortification:

The formulations in A, containing 100 mg of vitamin E acetate, was dispersed in a sample of each of the aforementioned beverages to form a 100 g stock solution of each beverage. 1.25 g, 2.50 g or 3.75 g of the stock solution of each beverage was placed into a beaker. Into the respective beakers were placed 28.75 g, 27.50 g or 26.25 g of each of the respective beverages. Each beaker was stirred to disperse the materials therein to form beverages containing 1.25 mg, 2.50 mg and 3.75 mg of vitamin E acetate per 30 g of beverage. Turbidity of the beverages was measured as described above.

The optical clarity results for the beverages of Examples 1 and 2 are shown in Table 1.

25

5

10

15

Table 1

Beverage	Examples	Ti	urbidity, NT	'U*
·	·	1.25 mg	2.50 mg	3.75 mg
Gatorade Lemon Ice	1	27	50	76
	2	5.6	6.8	7.3
Apple Juice	1	14.2	29	45
	2	2.2	2.3	3.4

* A NTU value of 10 or less is considered to be optically clear.

As can-be seen from Table 1, the beverages fortified with the formulation of the present invention were much less turbid than beverages fortified with the control formulation.

Example 3

A. Formulation:

Step 1. Vitamin E acetate (4 grams) and Polysorbate 80 (polyoxyethylene (20) sorbitan mono-oleate; 6 grams) were mixed in a suitable container and heated to about 40°C to give a homogeneous mixture.

Step 2. The above mixture was mixed with water in various proportions and stirred at 30-35°C for about 5 minutes or until the mixtures became gel-like. These mixtures were used for beverage fortification of Gatorade Lemon Ice.

B. Fortification:

The formulations in A, containing 100 mg of vitamin E acetate, were dispersed in a sample of the aforementioned beverage to form a 100 g stock solutions thereof. 1.25 g or 2.50 g of each of the stock solutions of the beverage were placed into a beaker. Into the respective beakers were placed 28.75 g or 27.50 g of the beverage. Each beaker was stirred to disperse the materials therein to form beverages containing 1.25 mg or 2.50 mg of vitamin E acetate per 30 g of beverage.

The results are shown in Table 2.

Table 2.

	Tu	rbidity, NTU*
Ratio of (vitamin E/polysorbate) to water	1.25 mg	2.50 mg
1/ 0.25	14.6	26
1/0.33	13.7	27
1/0.50	9.5	15.5

1/1	5.2	7.3
1/2	6.5	8.9
1/4	4.9	5.6
1/6	4.7	5.7
1/9	6.3	7.4
1/16	9.6	15.7
1/25	21	37
1/50	22	38
1/100	22	38
1/200	23	40
1/400	27	46

^{*} A value of NTU of 10 or less is considered to be optically clear.

Example 4

The stability of an inventive formulation, comprising 20 % vitamin E acetate, 30 % Tween 80 (polyoxyethylene (20) sorbitan mono-oleate), and 50 % water, was evaluated at room temperature (RT, ~ 22°C). The formulation was made in the same manner as other formulations of the present invetion as above and was placed into a suitable container for stability evaluation. The results of the stability evaluation are shown in Table 3 below.

10

5

Table 3

	% 1	Retention at RT*	
Lot#	Initial (IU/gram)	1.5 months	3 months
1	95.0 (195)	108 (216)	105 (209)
2	93.5 (187)	104 (208)	117 (235)

^{*} Based on the claim, which is 200 IU of vitamin E/gram

Example 5

15

20

The chemical and physical stability, as measured by percent vitamin E retention, presence or absence of ringing or precipitation, and optical clarity, of the formulation of Example 4 was evaluated in a variety of beverages. The results of the evaluation is shown in Table 4 below. The formulation was incorporated into the various beverages in a manner as described in Example 2 above.

Table 4

Beverage Type	Fortification Level (mg / 240			Vit. E Retention (%)	(etentic	(%) u			Ringing or precipitation after 6 mo storage	or ation no	Optical NTU	Optical Clarity* NTU
	.	Initial	1 Month	hth	3 Mc	3 Months	6 Months	nths)		Initial	9
					,							Months
			2°C	RT	၁့၄	RT	2°€	RT	2°C	RT	-	RT
Gatorade Lemon Ice	10	104	95	93	112	106	96	97	No	No	5.6	6.4
Gatorade Lemon Ice	20	104	66	101	104	100	101	101	No	No	7.6	8.6
Gatorade Lemon	10	116	107	105	119	121	111	110	No	8	11	114
Lime												
Gatorade Lemon	20	114	109	105	110	110	109	117	° N	°N	110	117
Lime												00,
Gatorade Orange	10	112	97	97	112	106	102	101	٥	٥ N	115	120
Gatorade Orange	20	105	86	66	103	104	97	103	No	S N	114	117
Lucky Leaf Apple	10	108	111	107	102	102	93	94	o Z	°Z	5.3 .3	*
Juice												1 1
Lucky Leaf Apple	20	107	104	104	101	101	94	96	o Z	o Z	8.7	:
Juice												

Initial (final) optical clarity: Lemon Ice: 4.7 (4.9); Lemon Lime: 108 (109); Orange: 113 (113) not determined Fortification Level is mg of vitamin E acetate per 240 ml of beverage.

The beverage or beverage concentrates supplemented with the optically clear composition of fat soluble vitamins of the present invention can be made by conventional means well known to those of ordinary skill in the art. In general, the optically clear composition of fat soluble vitamins can be incorporated into the beverage or beverage concentrates or syrups in the same manner in which the optically clear composition of fat soluble vitamins have been added in the above examples.

The beverage concentrates and syrups to which the optically clear composition of fat soluble vitamins have been added can be used to make final single strength beverage by blending the concentrate or syrup with an appropriate amount of water, usually about 1 part concentrate or syrup to abut 3 to 4 parts of water. In some instances, the water can either be carbonated or non-carbonated.

Examples of other beverages to which the optically clear composition of fat soluble vitamins can be added in nutritionally supplemental amounts include:

20

25

INSDOCID: <WO____

_9815195A1_I_>

5

10

- (a) "sparkling" orange juice comprising 55% orange juice and 45% carbonated water;
- (b) pear-grapefruit nectar comprising 25% pear juice, 20% grapefruit juice, the balance comprising 10% sucrose-water;
 - (c) kiwi-grapefruit drink comprising 20% kiwi fruit juice, 15% grapefruit juice, the balance comprising water;
- 30 (d) mixed fruit "cocktail" comprising 10% each of the juices of passion fruit, mango, guava, pineapple, papaya, banana, apricot, mandarin orange, pear and lime juices;
- (e) yogurt/fruit beverage comprising 20% milk products, 1% pectin, 20%
 pineapple juice, 10% shredded pineapple fruit pulp, 16% corn syrup, the balance comprising water;

- (f) cola beverage comprising 0.35% cola flavor emulsion, 11% sugar, 0.1% phosphoric acid, 0.1% citric and malic acids, caramel coloring, the balance comprising carbonated water;
- (g) full-strength orange juice;
- (h) full-strength apple juice;
- 10 (i) full-strength flavored cow's milk.

Claims:

- 1. An optically clear vitamin supplement comprising:
- 5 (A) (1) a fat soluble vitamin, and
 - (2) an emulsifier for (1), the ratio of (1) to (2) being from about 1:1 to about 1:19, and
 - (B) a liquid,

wherein the ratio of (A) to (B) is from about 1:0.5 to about 1:16.

10

- 2. A supplement according to claim 1 wherein (A)(1) is selected from the group consisting of vitamin E or its esters, vitamin A or its esters, vitamin K, and vitamin D3.
- 15 3. A supplement according to claim 2 wherein (A)(1) is vitamin E acetate.
 - 4. A supplement according to any one of claims 1-3, wherein (A)(2) is polyoxyethylene (20) sorbitan mono-oleate.
- 20 5. A supplement according to any one of claims 1-4, wherein (B) is water.
 - 6. A supplement according to any one of claims 1-5, where the ratio of (A)(1) to (A)(2) is from about 1:1 to about 1:9, preferably 1:1 to about 2:3, most preferably about 2:3.

- 7. A supplement according to any one of claims 1-6, wherein the ratio of (A) to (B) is from about 1:1 to about 1:9.
- 8. A nutritionally supplemented beverage comprising a mixture of
- 30 (a) at least about 0.05% by weight of fruit or cola flavor or at least about 3% by weight of fruit juice;
 - (b) an optical clear vitamin supplement according to any one of claims 1-7; and
 - (c) water if necessary to bring the beverage weight to 100%.
- 9. A beverage according to claim 8 wherein (b)(B) is water, the beverage to be supplemented, or mixtures thereof.

PCT/EP97/05421

- 10. A beverage according to claims 8 or 9 which additionally contains a sweetener.
- 11. A beverage according to claim 10 wherein the sweetener is selected from the group consisting of sucrose, fructose, glucose, high fructose corn syrup, corn syrup, and mixtures thereof.
- 12. A beverage according to claim 8 wherein the fruit juice is selected from the group consisting of grape, pear, passion fruit, pineapple, banana or banana puree, apricot, orange, lemon, grapefruit, apple, cranberry, tomato, mango, papaya, lime, tangerine, cherry, raspberry, carrot and mixtures thereof.

15

NSDOCID: <WO____9815195A1_I_>

INTERNATIONAL SEARCH REPORT

Inte .onal Application No

			PUI/EP 9/	/05421
A. CLASS IPC 6	FICATION OF SUBJECT MATTER A23L1/302 A23L1/035 A23L2	2/52 A61K9/00) A61K	9/107
According t	to International Patent Classification(IPC) or to both national da	ssification and IPC		
B. FIELDS	SEARCHED			
Minimum d IPC 6	ocumentation searched (classification system followed by class A23L A61K	ification symbols)		
Documenta	ation searched other than minimum documentation to the extent	that such documents are includ	ded in the fields se	arched
Electronic o	data base consulted during the international search (name of da	ita base and, where practical, s	search terms used))
		-		
	·			
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT			
Category °	Citation of document, with indication, where appropriate, of the	ne relevant passages		Relevant to claim No.
A	WO 95 24832 A (SMITHKLINE BEEC September 1995	CHAM) 21		1-12
	see claims see page 3, paragraph 1			
	see page 3, paragraph 3			
	see examples 3,4			
Α	WO 94 06310 A (SMITHKILNE BEEC March 1994			1-12
	see page 4, line 14-22; claims 3,5	; examples		
A,P	WO 97 10725 A (SMITHKILE BEECH	IAM) 27 March		1-12
	see claims			
	see page 3, line 29-36 see page 4, line 20-24			
Furt	her documents are listed in the continuation of box C.	χ Patent family m	embers are listed i	n annex.
° Special ca	ategories of cited documents :	"T" later document publi	shed after the inter	national filing date
consid	ent defining the general state of the art which is not dered to be of particular relevance	or priority date and cited to understand invention	not in conflict with	the application but
l filing o		"X" document of particul cannot be consider	ar relevance; the c	laimed invention
wnich	ant which may throw doubts on priority claim(s) or is cited to establish the publication date of another n or other special reason (as specified)	involve an inventive "Y" document of particul	step when the do	cument is taken alone
"O" docum	ent referring to an oral disclosure, use, exhibition or means	cannot be considered	ed to involve an inv ned with one or mo	ventive step when the ere other such docu-
"P" docum	ent published prior to the international filing date but han the priority date claimed	in the art. "&" document member o		us to a person skilled
Date of the	actual completion of theinternational search	Date of mailing of th		
2	4 February 1998	02/03/19	98	
Name and	mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2	Authorized officer		
	NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Fax: (+31-70) 340-3016	Van Moer	`, A	

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inte onal Application No PCT/EP 97/05421

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9524832 A	21-09-95	AU 1950395 A CA 2185584 A EP 0741523 A JP 9510106 T ZA 9502050 A	03-10-95 21-09-95 13-11-96 14-10-97 26-01-96
WO 9406310 A	31-03-94	AT 152584 T AU 671152 B AU 4979493 A CA 2144083 A CN 1087481 A DE 69310523 D DE 69310523 T EP 0660676 A JP 8504089 T MX 9305606 A NZ 255556 A US 5607707 A ZA 9306729 A	15-05-97 15-08-96 12-04-94 31-03-94 08-06-94 12-06-97 13-11-97 05-07-95 07-05-96 31-03-94 28-10-96 04-03-97 09-02-95
WO 9710725 A	27-03-97	AU 7086096 A	09-04-97